

Polyclonal antibody against Factor VIII-Related Antigen

Clone no. -

MONOSAN Ready To Use

---

|                           |   |
|---------------------------|---|
| Product name              | Polyclonal antibody against Factor VIII-Related Antigen           |
| Host                      | Rabbit  |
| Applications              | IHC-P   |
| Species reactivity        | human   |
| Conjugate                 | -   |
| Immunogen                 | Unknown or proprietary to MONOSAN and/or its suppliers            |
| Isotype                   | -   |
| Clonality                 | Monoclonal  |
| Clone number              | -   |
| Size                      | 7 ml  |
| Concentration             | n/a   |
| Format                    | -   |
| Storage buffer            | Tris Buffer, pH 7.3-7.7, containing 1% BSA and <0.1% Sodium Azide |
| Storage until expiry date | 2-8°C   |

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Polyclonal antibody against Factor VIII-Related Antigen

Clone no. -

MONOSAN Ready To Use

---

**Additional info**

Anti-Factor VIII-Related Antigen antibody reacts with endothelial cells and neoplastic blood cells. This antibody has helped to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi's sarcoma and cardiac myxoma. Not all endothelial cells synthesize (or store) this molecule; therefore, it should not be surprising that not all tumors of endothelial differentiation (benign or malignant) react with this antigen.

**References**

1. Nichols GE, et al. Am J Clin Pathol. 1992; 97:770-5
2. Falk S, et al. Am J Surg Pathol. 1993; 17:959-70
3. Meis-Kindblom JM, et al. Am J Surg Pathol. 1998; 22:683-97
4. Allison KH, et al. Am J Surg Pathol. 2004; 28:298-307
5. Peyvandi F, et al. Blood Transfus. 2011; 9 Suppl 2:s3-8

**FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES**