

Rabbit anti-Synaptophysin, clone MRQ-40

Clone no. MRQ-40

MONOSAN Ready To Use

---

|                           |   |
|---------------------------|---|
| Product name              | Rabbit anti-Synaptophysin, clone MRQ-40                           |
| Host                      | Rabbit  |
| Applications              | IHC-P   |
| Species reactivity        | human   |
| Conjugate                 | -   |
| Immunogen                 | Unknown or proprietary to MONOSAN and/or its suppliers            |
| Isotype                   | IgG   |
| Clonality                 | Monoclonal  |
| Clone number              | MRQ-40  |
| Size                      | 7 ml  |
| Concentration             | n/a   |
| Format                    | -   |
| Storage buffer            | Tris Buffer, pH 7.3-7.7, containing 1% BSA and <0.1% Sodium Azide |
| Storage until expiry date | 2-8°C   |

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Rabbit anti-Synaptophysin, clone MRQ-40

Clone no. MRQ-40

MONOSAN Ready To Use

---

**Additional info**

The antibody reacts with neuroendocrine cells of human adrenal medulla, carotid body, skin, pituitary, thyroid, lung, pancreas, and gastrointestinal mucosa. This antibody identifies normal neuroendocrine cells and neuroendocrine neoplasms. Diffuse, finely granular, cytoplasmic staining is observed, which probably correlates with the distribution of the antigen within neurosecretory vesicles. The expression of synaptophysin is independent of the presence of NSE or other neuroendocrine markers. Antisynaptophysin is an independent, broad-range marker of neural and neuroendocrine differentiation.

**References**

1. Wiedenmann, B, et al. Cell 1985;41:1017-1028
2. Navone, F et al. J Cell Biol 1986;103:2511-2527
3. Lyda MH et al. Hum Pathol. 2000 Aug;31(8):980-7
4. Skacel M et al. Appl Immunohistochem Mol Morphol. 2000 Sep;8(3):302-9
5. Morrison CD, Prayson RA. Semin Diagn Pathol. 2000 Aug;17(3):204-15

**FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES**