Product datasheet

MON1074



Mouse anti-CDw78, clone 60-3G2 (Monoclonal)Clone no.60-3G2

Product name	Mouse anti-CDw78, clone 60-3G2 (Monoclonal)		
Host	Mouse		
Applications	FC, IHC-fr, IF		
Species reactivity	human		
Conjugate	-		
Immunogen	In vitro activated B-cells		
lsotype	lgG1		
Clonality	Monoclonal		
Clone number	60-3G2		
Size	100 ug		
Concentration	100 ug/ml		
Format	-		
Storage buffer	PBS with 0.02% sodium azide		
Storage until expiry date	2-8°C		

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Product datasheet

MON1074

MONOSAN

Mouse anti-CDw78, clone 60-3G2 (Monoclonal) Clone no. 60-3G2

MONOSAN

Additional info

CDw78 (also called Ba antigen, Leu21 or LO panB a) is present on some immature and some mature B-cells. The antigen appears on B-cell progenitors preceding CD10, CD19, CD22, and CD37. It is expressed on resting B-cells and reappears and persists in the cytoplasm and on the cell surface until cytoplasmic Ig appears. Its expression is greatly increased after B-cell activation in vitro. It is also found on tissue macrophages and on epithelial cells, but not on T-cells, NK cells, monocytes, granulocytes, thymocytes or bone marrow stromal fibroblasts nor myeloid tissues. 60-3G2 was typed at CD workshop IV.

	~		
	tor	<u>' nn</u>	COC
1/C		CI	ces

1. Pinchouk VG. et al, Anticancer Research 8: 1377-1380 (1988)

- 2 Gluzman DF. et al, Tissue Antigens 33: 151 (1989)
- 3. Sidorenko SP. et al, Neoplasma 39: 3-9 (1992)

4. Moldenhauer et al, Leucocyte Typing IV, pp 155 162, (1989)

5. Pezzuto et al, Leucocyte Typing IV, pp 165 174, (1989)

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

www.monosan.com