

Mouse anti-INI-1, clone MRQ-27 (Monoclonal)

Clone no. MRQ-27

MONOSAN

Product name	Mouse anti-INI-1, clone MRQ-27 (Monoclonal)
Host	Mouse
Applications	IHC-P (1:100-1:500)
Species reactivity	human
Conjugate	-
Immunogen	Unknown or proprietary to MONOSAN and/or its suppliers
Isotype	IgG2a
Clonality	Monoclonal
Clone number	MRQ-27
Size	1 ml
Concentration	n/a
Format	-
Storage buffer	Tris Buffer, pH 7.3-7.7, containing 1% BSA and <0.1% Sodium Azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-INI-1, clone MRQ-27 (Monoclonal)

Clone no. MRQ-27

MONOSAN

Additional info

The INI-1 gene, which encodes a functionally uncharacterized protein component of the hSWI/SNF chromatin remodeling complex, is often mutated or deleted in malignant rhabdoid tumor (MRT). Two isoforms of INI-1 that differ by the variable inclusion of amino acids are potentially produced by differential RNA splicing. The morphology of MRTs can present challenges in differential diagnosis. The overall survival of MRTs relative to its potential mimics [medulloblastoma, supratentorial primitive neuroectodermal tumors (sPNETs)] is quite low, and thus differentiation from these other tumors is desirable. Lack of nuclear labeling by anti-INI-1 is characteristic of MRT. The majority of medulloblastomas and sPNETs are labeled by anti-INI-1. MRTs also originate from the kidney and soft tissues.

References

1. Bourdeaut F, et al. J Pathol. 2007; 211:323-30
2. Fowler DJ, et al. Fetal Pediatr Pathol. 2006; 25:159-68
3. Haberler C, et al. Am J SurgPathol. 2006; 30:1462-8
4. Janson K, et al. Pediatr Blood Cancer. 2006 Sep;47(3):279-84
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES