

Mouse anti-Adenovirus, clone 20/11 & 2/6 (Monoclonal)

Clone no. 20/11 & 2/6

MONOSAN

| | |
|---------------------------|---|
| Product name | Mouse anti-Adenovirus, clone 20/11 & 2/6 (Monoclonal) |
| Host | Mouse |
| Applications | IHC-P (1:25-1:100) |
| Species reactivity | human |
| Conjugate | - |
| Immunogen | Unknown or proprietary to MONOSAN and/or its suppliers |
| Isotype | IgG1-k |
| Clonality | Monoclonal |
| Clone number | 20/11 & 2/6 |
| Size | 1 ml |
| Concentration | n/a |
| Format | - |
| Storage buffer | Tris Buffer, pH 7.3-7.7, containing 1% BSA and <0.1% Sodium Azide |
| Storage until expiry date | 2-8°C |

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-Adenovirus, clone 20/11 & 2/6 (Monoclonal)

Clone no. 20/11 & 2/6

MONOSAN

Additional info

Adenovirus infection is associated with a broad spectrum of clinical disease in both children and adults. It has gained more attention as an important complication in patients who have undergone bone marrow or solid organ transplantation. The incidence of adenovirus infection in bone marrow transplant patients has been reported at 5-20%. Adenovirus infection on morphology should be differentially diagnosed from other virus infections, especially CMV infection. Anti-adenovirus can assist in this differential diagnosis by showing a round or crescent-shaped nuclear inclusion, generally within the surface epithelium and is exclusively intra-nuclear in location.

References

1. son, MG. Clin Infect Dis. 2006; 43: 331–9
2. Shayan K, et al. Arch Pathol Lab Med 2003;127:1615-8
3. -
4. -
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES