

Mouse anti-Alpha-Tubulin, clone TU-01 (Monoclonal)

Clone no. TU-01

MONOSAN

Product name	Mouse anti-Alpha-Tubulin, clone TU-01 (Monoclonal)
Host	Mouse
Applications	WB, IHC-P, ICC, IP, FC
Species reactivity	Mouse, Pig, Human, Turkey, Eisenia, Paramecium, Nicotiana, Yeast, Arabidopsis
Conjugate	-
Immunogen	Fraction of tubulin purified from porcine brain by two cycles of polymerization - depolymerization.
Isotype	IgG1
Clonality	Monoclonal
Clone number	TU-01
Size	0.1 mg
Concentration	1 mg/ml
Format	-
Storage buffer	Phosphate buffered saline (PBS) solution with 15 mM sodium azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-Alpha-Tubulin, clone TU-01 (Monoclonal)

Clone no. TU-01

MONOSAN

Additional info

The antibody TU-01 recognizes a defined epitope (aa 65-97) on N-terminal structural domain of alpha-tubulin. The microtubules are intracellular dynamic polymers made up of evolutionarily conserved polymorphic alpha/beta-tubulin heterodimers and a large number of microtubule-associated proteins (MAPs). The microtubules consist of 13 protofilaments and have an outer diameter 25 nm. Microtubules have their intrinsic polarity; highly dynamic plus ends and less dynamic minus ends. Microtubules are required for vital processes in eukaryotic cells including mitosis, meiosis, maintenance of cell shape and intracellular transport. Microtubules are also necessary for movement of cells by means of flagella and cilia. In mammalian tissue culture cells microtubules have their minus ends anchored in microtubule organizing centers (MTOCs). The GTP (guanosintriphosphate) molecule is an essential for tubulin heterodimer to associate with other heterodimers to form microtubule. In vivo, microtubule dynamics vary considerably. Microtubule polymerization is reversible and a populations of microtubules in cells are on their minus ends either growing or shortening – this phenomenon is called dynamic instability of microtubules. On a practical level, microtubules can easily be stabilized by the addition of non-hydrolysable analogues of GTP (eg. GMPPCP) or more commonly by anti-cancer drugs such as Taxol. Taxol stabilizes microtubules at room temperature for many hours. Using limited proteolysis by enzymes both tubulin subunits can be divided into N-terminal and C-terminal structural domains. The alpha-tubulin (relative molecular weight around 50 kDa) is globular protein that exists in cells as part of soluble alpha/beta-tubulin dimer or it is polymerized into microtubules. In different species it is coded by multiple tubulin genes that form tubulin classes (in human 6 genes). Expressed tubulin genes are named tubulin isotypes. Some of the tubulin isotypes are expressed ubiquitously, while some have more restricted tissue expression. Alpha-tubulin is also subject of numerous post-translational modifications. Tubulin isotypes and their posttranslational modifications are

References

1. Viklicky V, et al. Cell Biol Int Rep. 1982 Aug;6(8):725-31
2. Grimm M, et al. Biochim Biophys Acta. 1987 Jul 24;914(1):83-8.
3. Linhartova I, et al. Biochem J. 1992 Dec 15;288 (Pt 3):919-24.
4. Draber P, et al. Eur J Cell Biol. 1986 Jun;41(1):82-8
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES