

Mouse anti-CNF1, clone NG8 (Monoclonal)

Clone no. NG8

MONOSAN

Product name	Mouse anti-CNF1, clone NG8 (Monoclonal)
Host	Mouse
Applications	FUNC,ELISA,WB
Species reactivity	n/a
Conjugate	-
Immunogen	Unknown or proprietary to MONOSAN and/or its suppliers
Isotype	IgG2a
Clonality	Monoclonal
Clone number	NG8
Size	1 ml
Concentration	100 ug/ ml
Format	-
Storage buffer	PBS with 0.1% BSA and 0.02% sodium azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-CNF1, clone NG8 (Monoclonal)

Clone no. NG8

MONOSAN

Additional info

The monoclonal antibody NG8 is specific for Cytotoxic necrotizing factor type 1 (CNF1) of uropathogenic Escherichia coli. CNF1 and CNF2 belong to a family of bacterial toxins that target the small GTP-binding Rho proteins that regulate the actin cytoskeleton. Members of this toxin family typically inactivate Rho; however, CNF1 and the highly related CNF2 activate Rho by deamidation. CNF1 is more frequently associated with E.coli strains that cause extraintestinal infections in humans, particularly those of the urinary tract (such as cystitis, pyelonephritis and prostatitis). In CNF1-producing uropathogenic E. coli strains, CNF1 is chromosomally encoded and typically resides on a pathogenicity island that also contains hemolysin and P fimbria-related genes. Both CNF1 and the highly related, plasmid-encoded CNF2 are monomeric, cytoplasmic toxins of approximately 115 kDa. CNF1 can be structurally organized into three functional domains the N-terminal, central and the C-terminal domain. The latter exhibits the catalytic activity of the toxin. Monoclonal antibody NG8 recognizes an epitope between amino acids 704 and 730 of the C-terminal enzymatic domain. NG8 specifically neutralizes CNF1 while lacking activity for CNF2.

References

1. Meysick; K et al. Infect Immun 2001; 69: 2066
2. McNichol, B et al Infect Immun 2007, 75: 5095
3. -
4. -
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES