

Mouse anti-CD30, clone MEM-268 (Monoclonal)

Clone no. MEM-268

MONOSAN

---

Product name	Mouse anti-CD30, clone MEM-268 (Monoclonal)
Host	Mouse
Applications	FC , IHC-P
Species reactivity	Human
Conjugate	-
Immunogen	Expression vector containing CD30 cDNA (booster suspension of THP-1 cell line)
Isotype	IgG
Clonality	Monoclonal
Clone number	MEM-268
Size	0.1 mg
Concentration	1 mg/ml
Format	-
Storage buffer	Phosphate buffered saline (PBS) solution with 15 mM sodium azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

## Mouse anti-CD30, clone MEM-268 (Monoclonal)

Clone no. MEM-268

MONOSAN

**Additional info**

CD30 is a type I transmembrane glycoprotein of the TNF receptor superfamily. CD30 was originally identified as a cell surface antigen of Hodgkins and Reed-Sternberg cells using monoclonal antibody Ki-1. The ligand for CD30 is CD30L (CD153). The binding of CD30 to CD30L mediates pleiotropic effects including cell proliferation, activation, differentiation, and apoptotic cell death. CD30 has a critical role in the pathophysiology of Hodgkin's disease and other CD30+ lymphomas. CD30 acts as a costimulatory molecule in thymic negative selection. In addition to its expression on Hodgkin's and Reed-Sternberg cells, CD30 is also found in some non-Hodgkin's lymphomas (including Burkitt's lymphomas), virus-infected T and B cells, and on normal T and B cells after activation. In T cells, CD30 expression is present on a subset of T cells that produce Th2-type cytokines and on CD4+/CD8+ thymocytes that co-express CD45RO and the IL4 receptor. Soluble form of CD30 (sCD30) serves as a marker reflecting Th2 immune response.

**References**

1. -
2. -
3. -
4. -
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES