

Mouse anti-MHC II DR (alpha + beta chain), clone Bra30 (Monoclonal)

Clone no. Bra30

MONOSAN

---

Product name	Mouse anti-MHC II DR (alpha + beta chain), clone Bra30 (Monoclonal)
Host	Mouse
Applications	FC, IHC-fr, IHC-P
Species reactivity	human
Conjugate	-
Immunogen	REH (human ALL cell line)
Isotype	IgG2a-K
Clonality	Monoclonal
Clone number	Bra30
Size	100 ug
Concentration	100 ug/ml
Format	-
Storage buffer	PBS with 0.02% sodium azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-MHC II DR (alpha + beta chain), clone Bra30 (Monoclonal)

Clone no. Bra30

MONOSAN

**Additional info**

MHC class II molecules are encoded by polymorphic MHC genes and consist of a non-covalent complex of an  $\alpha$  and  $\beta$  chain. Helper T lymphocytes bind antigenic peptides presented by MHC class II molecules. MHC class II molecules bind 13-18 amino acid antigenic peptides. Accumulating in endosomal/lysosomal compartments and on the surface of B cells, HLA-DM and -DO molecules regulate binding of exogenous peptides to class II molecules (HLA-DR) by sustaining a conformation that favors peptide exchange. The differential structural properties of MHC class I and class II molecules account for their respective roles in activating different populations of T lymphocytes.

**References**

1. Chorvath B et al. Neoplasma 34(4): 417-425 (1987)
2. Horejsi V et al. Tissue Antigens 32(1): 6-11 (1988)
3. Polakova K et al. Neoplasma 32(6): 641-8 (1985)
4. -
5. -

**FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES**