

Mouse anti-CDw78/MHC II DR, clone 60-3G2 (Monoclonal)

Clone no. 60-3G2

MONOSAN

---

Product name	Mouse anti-CDw78/MHC II DR, clone 60-3G2 (Monoclonal)
Host	Mouse
Applications	FC, IHC-fr, IF
Species reactivity	human
Conjugate	-
Immunogen	human B-cell line Daudi
Isotype	IgG1-K
Clonality	Monoclonal
Clone number	60-3G2
Size	100 ug
Concentration	100 ug/ml
Format	-
Storage buffer	PBS with 0.02% sodium azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-CDw78/MHC II DR, clone 60-3G2 (Monoclonal)

Clone no. 60-3G2

MONOSAN

**Additional info**

CDw78 (also called Ba antigen, Leu21 or LO panB a) is present on some immature and some mature B-cells. The antigen appears on B-cell progenitors preceding CD10, CD19, CD22, and CD37. It is expressed on resting B-cells and reappears and persists in the cytoplasm and on the cell surface until cytoplasmic Ig appears. Its expression is greatly increased after B-cell activation in vitro. It is also found on tissue macrophages and on epithelial cells, but not on T-cells, NK cells, monocytes, granulocytes, thymocytes or bone marrow stromal fibroblasts nor myeloid tissues. 60-3G2 was typed at CD workshop IV.

**References**

1. Pinchouk VG. et al, Anticancer Research 8: 1377-1380 (1988)
2. Gluzman DF. et al, Tissue Antigens 33: 151 (1989)
3. Sidorenko SP. et al, Neoplasma 39: 3-9 (1992)
4. Moldenhauer et al, Leucocyte Typing IV, pp 155 162, (1989)
5. Pezzuto et al, Leucocyte Typing IV, pp 165 174, (1989).

**FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES**