

# Recombinant Mouse Interleukin 18 (without BSA)

**Code No.****Quantity****Form****B004-2****200 µg x 1 vial****lyophilized**

**BACKGROUND:** Interleukin 18 (IL-18) is a 18 kDa novel cytokine which identified as a costimulatory factor for production of interferon- $\gamma$  (IFN- $\gamma$ ) in response to toxic shock and shares functional similarities with IL-12. IL-18 is synthesized as a precursor 24 kDa molecule without a signal peptide and must be cleaved to produce an active molecule. IL-1 converting enzyme (ICE, Caspase-1) cleaves pro-IL-18 at aspartic acid in the P1 position, producing the mature, bioactive peptide that is readily released from the cells. It is reported that IL-18 is produced from Kupffer cells, activated macrophages, keratinocytes, intestinal epithelial cells, osteoblasts, adrenal cortex cells and murine diencephalon.

IFN- $\gamma$  is produced by activated T or NK cells and plays critical roles in the defense against microbial pathogens. IFN- $\gamma$  activates macrophages, enhances NK activity and B cell maturation, proliferation and Ig secretion, induces MHC class I and II antigens, and inhibits osteoclast activation.

IL-18 acts on T helper 1-type T cells and in combination with IL-12 strongly induces them to produce IFN- $\gamma$ . Pleiotropic effects of IL-18 has also been reported, such as, enhancement production of IFN- $\gamma$  and GM-CSF in peripheral blood mononuclear cells, production of T helper type 1 cytokines, IL-2, GM-CSF and IFN- $\gamma$  in T cells, enhancement of Fas ligand expression by T helper type 1 (Th1) cells.

**DESCRIPTION :** cDNA encoding the matured Mouse IL-18 protein sequence (corresponding to a.a. 36-192) was expressed in *E. coli*.

**PURITY :** Greater than 90% purity as confirmed on SDS-PAGE by Coomassie brilliant blue staining.

**MOLECULAR WEIGHT:** 18kDa

**ENDOTOXIN LEVEL:** Less than 0.1 ng per 1 µg of recombinant IL-18 protein, measured by LAL method.

**SUPPLIED :** Lyophilized in PBS containing 1 % sucrose. Reconstitute in 2 ml of ice-cold distilled water on ice.

**STORAGE :** Lyophilized IL-18 is stable for 24 months from the date of manufacture when stored at -20°C or below. After reconstitution, avoid repeated freeze/thaw cycles. The IL-18 can be stored for 1 week at 4°C. For storage, prepare appropriate aliquots and freeze them at -80°C.

**ACTIVITY :** Induction of IFN- $\gamma$  by Mouse IL-18 receptor transfected KG-1 cell (KG-1 cell: Human myelomonocyte; ATCC CCL246) in response to the Mouse IL-18 was measured by Human IFN- $\gamma$  ELISA.

The activity of lot 010 is as follow;

IL-18 conc. (ng/mL)	IFN- $\gamma$ induction (IU/mL)*
20	180.3
40	264.6

\*IFN- $\gamma$  producing activity of the sample cells can vary depending on cell conditions.

**IFN- $\gamma$  PRODUCTION ASSAY :**

- 1) KG-1 transfectant cells were cultured at  $3 \times 10^6$  cells/mL for two days at 37°C in 5% CO<sub>2</sub> incubator with RPMI 1640.
- 2) After two days of preculture, the cell concentration was adjusted to  $2 \times 10^6$  cells/mL and incubated for 46-48 hours at 37°C in 5% CO<sub>2</sub> incubator with RPMI 1640 in the presence of IL-18.
- 3) The culture supernatant were recovered and the amount of IFN- $\gamma$  were measured by Human IFN- $\gamma$  ELISA (code# IM1743: MBL).

**RESEARCH USE :** For research use only. Not for use in *in vitro* diagnostic procedures for clinical diagnosis.

**REFERENCE:**

- 1) Okamura H., et al. *Nature* **378**, 88-91 (1995)
- 2) Ushio S., et al. *J. Immunol.* **156**, 4274-4279 (1996)
- 3) Micallef M., et al. *Eur. J. Immunol.* **26**, 1647-1651 (1996)
- 4) Tao D, et al. *Cell Immunol.* **173**, 230-235 (1998)

B004-2

October 9, 2003

**Medical & Biological Laboratories Co., Ltd.**

15 B Constitution Way, Woburn, MA 01801

TEL : (781) 939-6964 FAX : (781) 939-6963 www.mblintl.com